

IOWA STATE UNIVERSITY

Digital Repository

Animal Science Publications

Animal Science

2000

Rapid communication: Porcine vitamin D-25-hydroxylase maps to chromosome 5

Daniel C. Ciobanu

Iowa State University

R. K. Gill

Medical University of South Carolina

Max F. Rothschild

Iowa State University, mfrothsc@iastate.edu

N. H. Bell

Medical University of South Carolina

Follow this and additional works at: http://lib.dr.iastate.edu/ans_pubs



Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), and the [Genetics and Genomics Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/ans_pubs/276. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Rapid communication: Porcine vitamin D-25-hydroxylase maps to chromosome 5

Abstract

Locus Name. Vitamin D-25-hydroxylase. Species. *Sus scrofa*. Locus Symbol. CYP25. Source of Primers. PCR primers were designed from the published sequence for porcine vitamin D-25-hydroxylase (CYP25) cDNA (GenBank accession no. Y16417). The position of the forward and reverse primers corresponds to 506 to 530 bp and 830 to 854 bp, respectively, of the cDNA. The primer pair amplified a 348-bp fragment of the cDNA and an 862-bp fragment from the pig genomic DNA. The 862-bp fragment (Gen- Bank accession no. AF284378) was subcloned into a pNoTA vector and identity was confirmed by sequencing. Genomic sequences from 1 to 74, 169 to 329, and 749 to 862 showed 100% homology to the cDNA.

Keywords

Gene Mapping, Oxygenases, Pigs, Vitamin D

Disciplines

Agriculture | Animal Sciences | Genetics and Genomics

Comments

This is an article from *Journal of Animal Science* 78 (2000): 3193, doi:/2000.78123193x. Posted with permission.

Rapid communication: Porcine vitamin D-25-hydroxylase maps to chromosome 5¹

D. C. Ciobanu*, R. K. Gill†, M. F. Rothschild*, and N. H. Bell†²

*Department of Animal Science, Iowa State University, Ames 50011 and

†Department of Medicine, Medical University of South Carolina, Charleston 29425

Locus Name. Vitamin D-25-hydroxylase.

Species. *Sus scrofa*.

Locus Symbol. *CYP25*.

Source of Primers. PCR primers were designed from the published sequence for porcine vitamin D-25-hydroxylase (*CYP25*) cDNA (GenBank accession no. Y16417). The position of the forward and reverse primers corresponds to 506 to 530 bp and 830 to 854 bp, respectively, of the cDNA. The primer pair amplified a 348-bp fragment of the cDNA and an 862-bp fragment from the pig genomic DNA. The 862-bp fragment (GenBank accession no. AF284378) was subcloned into a pNoTA vector and identity was confirmed by sequencing. Genomic sequences from 1 to 74, 169 to 329, and 749 to 862 showed 100% homology to the cDNA.

Primer Sequences. Primers derived from the porcine sequences were as follows: forward, 5'-GGG CAA GAA GGG GTC GTT GGA GGA GTC-3', and reverse, 5'-ACT CGA CTA GTG TCT CGT GTT CTA C-3'.

Method of Detection. PCR was performed in 50- μ L reactions that included 1 μ L *AdvanTaq* (solution diluted 1:50) DNA polymerase (Clontech, Palo Alto, CA), 16 mM (NH)₄SO₄, 67.5 mM Tris buffer pH 8.8, 100 μ M dNTPs, 1.5 mM MgCl₂, 0.01% Tween-20, and 50 ng of genomic DNA. PCR was carried out in a Perkin Elmer GeneAmp 9600 PCR System (Norwalk, CT). The cycling conditions included initial denaturation at 95°C for 3 min followed by 30 cycles at 95°C for 45 s, 60°C for 1 min, 72°C for 2 min, and final extension at 72°C for 10 min. From the cDNA information it can be deduced that the 862-bp fragment of the porcine *CYP25* gene produced by PCR contained two introns and portions of two exons and a full exon.

Chromosome Location. The 862-bp fragment of the porcine *CYP25* gene was used for physical mapping by using the INRA-University of Minnesota porcine Radia-

tion Hybrid panel (Hawken et al., 1999). LOD values of 2.5 or greater obtained by the two-point analysis procedure are presented in Table 1; values of 4.8 were considered significant evidence for linkage. The results showed that the *CYP25* gene locus is located on the end of the short arm of pig chromosome 5 near the aconitase gene locus (*ACO2*). The retention fraction was 34%. Based on previous physical mapping results for *ACO2*, the position of the pig *CYP25* gene is likely on the chromosome 5p14–15 region.

Comments. A porcine vitamin D-25-hydroxylase previously was cloned and characterized (Postlind et al., 1997). The gene codes for a cytochrome P450 microsomal enzyme that is expressed in liver and kidney. Porcine *CYP25* cDNA consists of 1,652 bp, and the purified protein consists of 500 amino acids and catalyzes equally the conversion of vitamin D₂ and vitamin D₃ to 25-hydroxyvitamin D₂ and 25-hydroxyvitamin D₃, respectively (Axen et al., 1994; Postlind et al., 1997). Porcine *CYP25* therefore was proposed to be the true vitamin D-25-hydroxylase. Porcine *CYP25* is 83% homologous with human *CYP2D6*, an enzyme involved in the metabolism of debrisoquine and other drugs (Postlind et al., 1997). No *CYP2D6* activity was detected in pig liver (Skaanild and Friis, 1999). As indicated in the present report, the porcine *CYP25* gene locus maps to pig chromosome 5p14–15, close to the *ACO2* gene locus (Yasue et al., 1999). This region is syntenic to human chromosome 22q12–13 containing the loci for the human *CYP2D6* and *ACO2* genes (Gonzalez et al., 1988; Gough et al., 1993). The location of the *CYP25* gene in the RH map is in a telomeric region with few markers mapped, and this makes it difficult to ascertain the exact position of the gene. The biological relationship, if any, between porcine *CYP25* and human *CYP2D6* remains to be determined.

Literature Cited

- Axen, E., T. Bergman, and K. Wikvall. 1994. Microsomal 25-hydroxylation of vitamin D₂ and vitamin D₃ in pig liver. *J. Steroid Biochem. Mol. Biol.* 51:97–106.
- Gonzalez, F. J., F. Vilbois, J. P. Hardwick, O. W. McBride, D. W. Nebert, H. V. Gelboin, and U. A. Meyer. 1988. Human debrisoquine 4-hydroxylase (P450IID1): cDNA and deduced amino acid sequence and assignment of the *CYP2D* locus to chromosome 22. *Genomics* 2:174–179.
- Gough, A. C., C. A. D. Smith, S. M. Howell, C. R. Wolf, S. P. Bryant, and N. K. Spurr. 1993. Localization of the *CYP2D* gene locus to

¹This work was financially supported in part by PIC group and the Iowa Agric. and Home Econ. Exp. Sta., Ames, paper no. J-18858, project no. 3148, by Hatch Act and State of Iowa funds, and by an institutional grant from the Medical University of South Carolina.

²Correspondence: 114 Doughty Street, P.O. Box 250775 (phone: (843) 876-5162; fax: (843) 876-5163; E-mail: belln@musc.edu).

Received May 8, 2000.

Accepted August 3, 2000.

©2000 American Society of Animal Science. All rights reserved.

Table 1. Mapping the *CYP25* gene with the INRA-University of Minnesota porcine Radiation Hybrid panel: two-point analysis^a

Order	Chromosome	Marker	--	- +	+ -	++	Distance (ray)	LOD ^b
1	5	<i>ACO2</i>	72	4	19	21	0.64	6.17
2	5	<i>SW1482</i>	67	9	22	18	1.00	2.93
3	5	<i>DK</i>	52	25	13	28	1.03	2.77

^a+ +, Hybrids in which both markers were detected; --, Hybrids in which both markers were not detected; - + and + -, Hybrids in which one of the markers was detected and one not.
^bLOD values of 4.8 or greater are considered significant.

human chromosome 22q13.1 by polymerase chain reaction, in situ hybridization and linkage analysis. *Genomics* 15:430–432.

Hawken, R. J., J. Murtaugh, G. H. Flickinger, M. Yerle, A. Robic, D. Milan, J. Gellin, C. W. Beattie, L. B. Schook, and L. J. Alexander. 1999. A first-generation porcine whole-genome radiation hybrid map. *Mamm. Genome* 10:824–830.

Postlind, H., E. Axen, T. Bergman, and K. Wikvall. 1997. Cloning, structure and expression of a cDNA encoding vitamin D₃ 25-hydroxylase. *Biochem. Biophys. Res. Commun.* 241:491–497.

Skaanild, M. T., and C. Friis. 1999. Cytochrome P450 sex differences in minipigs and conventional pigs. *Pharmacol. Toxicol.* 85:174–180.

Yasue H., N. Hisamatsu, T. Awata, Y. Wada, and H. Kusumoto. 1999. Clarification of the order of acrosin and aconitase 2 genes on the physical and linkage maps of porcine chromosome 5. *Anim. Genet.* 30:161–162, 1999.

Key Words: Gene Mapping, Oxygenases, Pigs, Vitamin D

J. Anim. Sci. 2000. 78:3193–3194